

Fulde-Ferrell-Larkin-Ovchinnikov states in vortex lattices

Predrag Nikolić

Department of Physics, Rice University, Houston, TX 77005, USA

(Dated: April 15, 2009)

Abstract

The magnetism of superconductors has not been fully explored despite many years of research and its great significance to both theory and applications. Here we present a unified perspective on how quantum electromagnetic vortices and electron-spin textures join to protect superconductivity against the destructive power of large magnetic fields. Using rapidly rotating ultra-cold gasses of fermionic atoms as a model system amenable to detailed experimental exploration, we discover a hierarchy of Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) phases in which a metal or a band-insulator of unpaired particles coexists with a spatially modulated superfluid. Quantum fluctuations can transform these phases into strongly correlated vortex liquid metals or insulators. We elucidate a general criterion for the stability of FFLO phases and discuss implications for the quantum Hall and vortex liquid phenomena in the pseudogap state of high temperature superconductors.

Some fundamental aspects of magnetism in superconductors remain unclear despite many years of research. Several mechanisms for the survival of superconductivity in large magnetic fields have been studied: vortex lattice formation [1], re-entrant superconductivity [2, 3, 4], and the relaxation of Zeeman effect through spontaneous development of spatial modulations (FFLO states) [5, 6]. Only the first mechanism has been unambiguously observed in nature [7]. Fortunately, a new frontier has been opened in atomic physics to observe and study these phenomena. Here we analyze in detail the very rich phase diagram of rotating fermionic atoms with attractive interactions near unitarity in two dimensions.

Electronic systems are very similar to trapped ultra-cold fermionic alkali atoms [8]. The orbital effect of magnetic field can be created in atomic systems by rotation. Coriolis forces in the rotating frame implement a uniform flux density which can produce quantized vortices and quantum Hall effect. The paramagnetic or Zeeman effect is created by preparing a system with unequal number of atoms in the two lowest-energy hyperfine states, which essentially have the same dynamics as electron spin and will be referred to as “spin” from now on. The Zeeman and orbital effects can be independently controlled. The strength of attractive interactions between atoms, responsible for superfluidity, is easily tuned using Feshbach resonances. Therefore, cold atoms are ideal systems for exploring a wide range of phenomena related to superconductivity.

The competition between superfluidity and Zeeman effect is normally resolved by forming either an unpolarized superfluid of Cooper pairs, or a spin-polarized normal state [9, 10, 11, 12]. The elusive FFLO state is a compromise in which the superfluid and spin-polarization coexist but alternate in space in a periodic pattern. FFLO states have been independently studied in condensed matter and high energy physics [13]. Three-dimensional [14, 15] and two-dimensional [16, 17, 18, 19, 20, 21] FFLO states have been considered in superconductors and heavy-fermion metals. The latter studies were motivated by the layered structure of materials, which allows controlling independently the orbital and Zeeman effects by tilting the magnetic field with respect to the layers. In order to directly address the role of these effects in stabilizing two-dimensional FFLO states, we depart from the earlier studies and focus on the lowest temperatures where Landau level quantization is important. This will also establish a connection between re-entrant superfluidity, FFLO phenomena and strongly correlated insulators of general interest in unconventional superconductivity, using a previously unexplored rotating cold-atom perspective (for a review of past research,

mostly on rotating bosons or three-dimensional fermionic systems, see [22] and references therein).

Here we discover a sequence of FFLO phases at zero temperature in which a metal or a band insulator of unpaired atoms coexists with a superfluid hosting a vortex lattice. Surrounding these phases are strongly correlated vortex liquid states and atomic quantum Hall insulators. Our results clarify the most general stabilizing mechanism for FFLO states and provide a unified perspective on the fate of superconductors in high magnetic fields. We also elucidate the potential of trapped rotating ultra-cold atoms to be the first system that clearly realizes the FFLO states. This and other studies in the atomic physics setting are likely to shed light on the possible role of FFLO states in superconductors, heavy-fermion metals, quantum chromodynamics and astrophysics. Finally, our results point to experimental circumstances for the cold-atom realization and exploration of strongly correlated states which shape the unconventional properties of high-temperature superconductors.

The quantitative applicability of the following universal theory to ultra-cold fermionic atoms near the Feshbach resonance stems from the unitarity limit. The scattering length of two-body interactions diverges at the resonance, so that the microscopic details of interactions become irrelevant and all properties of the many-body system become functions of only dimensionless ratios of externally controlled parameters. Such scaling laws are protected inside all superfluid phases, but may break down due to rotation in two dimensions when strongly correlated vortex liquid states are formed [23].

We consider a Bogoliubov-de Gennes (BdG) model in grand-canonical ensemble which arises as a saddle-point approximation in the quantum field theory of neutral fermionic particles near unitarity [23]. This approximation becomes quantitatively accurate at sufficiently large densities or sufficiently far from the Feshbach resonance in the BCS (Bardeen, Cooper, Schrieffer) limit, but remains extremely useful even in the numerically accessible regimes for elucidating various qualitative features which may be observed in experiments. Rotation at angular velocity ω is described by a static gauge field $\mathbf{A}$ in the rotating frame which satisfies $\nabla \times \mathbf{A} = 2m\omega\hat{\mathbf{z}}$, where m is the mass of atoms. We assume that the centrifugal forces are almost exactly cancelled by the potential which traps atoms. The Zeeman effect occurs in finite Zeeman field h which together with the global chemical potential μ sets the chemical potentials for the two hyperfine species, $\mu_{\uparrow} = \mu + h$ and $\mu_{\downarrow} = \mu - h$.

In the absence of interactions all atoms would be localized in their Landau orbitals. Such

localization facilitates Cooper pairing and allows superfluidity to occur even in the deep quantum limit at high rotation rates [2, 3, 4]. The saddle-point approximation regards the superfluid as a macroscopic condensate of Cooper pairs into a single state given by the order parameter function:

$$\Phi(\mathbf{r}) = \sum_n \sum_{l=0}^{n_v-1} \phi_{n,l} \sum_j \Phi_n \left(\mathbf{r} \sqrt{\frac{4m\omega}{\hbar}} ; \frac{(l + n_v j)\delta q}{\sqrt{4m\hbar\omega}} \right). \quad (1)$$

The amplitudes $\phi_{n,l}$ describe the occupation of bosonic Landau levels $n \in \{0, 1, 2, \dots\}$ at wavevectors $q_x = \text{integer} \times \delta q$ in x -direction. The Landau level wavefunctions of dimensionless coordinates $\vec{\xi} = (\xi_x, \xi_y)$ are:

$$\Phi_n(\vec{\xi}; \eta) = \frac{1}{\sqrt{2^n n!}} e^{in\xi_x} e^{-\frac{1}{2}(\xi_y + \eta)^2} H_n(\xi_y + \eta) \quad (2)$$

in Landau gauge, where H_n are Hermite polynomials. This realizes a vortex lattice with periods $\Delta x = 2\pi\hbar/\delta q$ and $\Delta y = n_v \delta q (4m\omega)^{-1}$ containing n_v vortices per unit-cell. The hexagonal Abrikosov lattice has $n_v = 2$, $\delta q = (4\pi\sqrt{3}m\hbar\omega)^{1/2}$, and $\phi_{n,1} = (-1)^n \phi_{n,0}$.

The BdG Schrödinger equation determines the quantum-mechanical spectrum of fermionic quasiparticle excitations in the condensate:

$$\begin{pmatrix} \frac{[-i\hbar\nabla - \mathbf{A}(\mathbf{r})]^2}{2m} - \mu - h & \Phi(\mathbf{r}) \\ \Phi^*(\mathbf{r}) & -\frac{[-i\hbar\nabla + \mathbf{A}(\mathbf{r})]^2}{2m} + \mu - h \end{pmatrix} \begin{pmatrix} u_{\mathbf{n}}(\mathbf{r}) \\ v_{\mathbf{n}}(\mathbf{r}) \end{pmatrix} = E_{\mathbf{n}}(\phi_{n,l}) \begin{pmatrix} u_{\mathbf{n}}(\mathbf{r}) \\ v_{\mathbf{n}}(\mathbf{r}) \end{pmatrix}. \quad (3)$$

Here, $u_{\mathbf{n}}$ and $v_{\mathbf{n}}$ are the particle and hole wavefunctions of opposite spins in the eigenstate with quantum numbers $\mathbf{n}$. We solve this equation in the basis of Landau orbitals using exact diagonalization. The energy spectrum $E_{\mathbf{n}}$ determines the free energy $\mathcal{F}(\phi_{n,l})$ in the saddle-point approximation [24]:

$$\begin{aligned} \mathcal{F}(\phi_{n,l}) = & - \sum_{\mathbf{n}} \left[k_B T \ln (1 + e^{-\beta(E_{\mathbf{n}}(\phi_{n,l}) + h)}) + k_B T \ln (1 + e^{-\beta(E_{\mathbf{n}}(\phi_{n,l}) - h)}) \right. \\ & \left. + E_{\mathbf{n}}(\phi_{n,l}) - E_{\mathbf{n}}(0) \right] + \left[\frac{m\nu}{4\pi\hbar^2} - \Pi_{0,0}(0,0) \Big|_{\mu=h=T=0} \right] \int d^2r |\Phi(\mathbf{r})|^2. \end{aligned} \quad (4)$$

Here, $\beta = (k_B T)^{-1}$, k_B is Boltzmann constant, and $\nu = -a_z/a$ measures the strength of attractive interactions between particles through the two-body scattering length a and the confinement scale a_z in z -direction which renders the system two-dimensional. Therefore, ν is proportional to the detuning from the Feshbach resonance in experiments. The ultra-violet behavior is regulated by rotation, and the subtracted pairing susceptibility $\Pi_{n_1, n_2}(i\omega, p_x)$

assures that $\nu = 0$ corresponds to unitarity [23]. The global minimum of $\mathcal{F}(\phi_{n,l})$ determines $\phi_{n,l}$ in the thermodynamic equilibrium. Note that this approach is not limited to the vicinity of second order transitions like the earlier studies.

The BdG spectrum $E_{\mathbf{n}}$ in the superfluid at $h = 0$ has a particle-hole symmetric band-structure reminiscent of broadened Landau levels at high energies. The low energy bands can be associated to quantum tunnelling of quasiparticles between vortex cores. The Zeeman field h acts as a chemical potential for the BdG quasiparticles which breaks the particle-hole symmetry. At zero temperature all states with $E_{\mathbf{n}} < h$ are occupied, and with $E_{\mathbf{n}} > h$ are empty. There is a “metal-insulator” quantum phase transition each time h crosses a band-edge [25]. As long as superfluidity survives, the extra quasiparticles at energies $0 < E_{\mathbf{n}} < h$ carry spin polarization which is mostly concentrated near vortex cores. Translational and rotational symmetries are spontaneously broken by both the superfluid density (vortex lattice) and the coexisting spin-polarization, therefore these states are FFLO states.

In the traditional FFLO picture the order parameter spontaneously develops spatial modulations in response to Zeeman effect alone. If the lowest positive energy in the resulting quasiparticle band-structure is larger than the pairing gap, then the FFLO state cannot be stable because it would breach the Pauli-Clogston limit, making it more advantageous to align all spins with the Zeeman field than to form singlet Cooper pairs. This problem is avoided in the presence of a vortex lattice. The energy gap for the vortex core states is always lower than the pairing gap (see Fig.1).

We numerically minimized the free energy density (4) using a fermionic basis of 72 quasi-periodic states in each of the 50 lowest Landau levels. The global minimum was calculated among order parameters within the lowest five bosonic Landau levels and $n_v = 2$, $\delta q = (4\pi\sqrt{3}m\hbar\omega)^{1/2}$, allowing a variety of lattice structures. All results are summarized by the zero-temperature phase diagram in Fig.2.

The superfluid is divided into a number of phases with either gradually varying polarization, or polarization saturated at an integer number of spins per vortex. The unpaired polarized atoms form a Fermi surface in the former case, and a band-insulator in the latter case. At least some of the “metal-insulator” transitions are first order in terms of the superfluid order parameter, which allows them to persist at finite temperatures despite the loss of the sharp distinction between “metals” and “insulators” of the vortex core fermions. The phase diagram contains a sequence of quantum tri-critical points where the transitions

between atomic quantum Hall insulators meet the first and second order superfluid-insulator transitions. The superfluid boundary retains its pair-breaking first order transitions below some finite temperatures.

Interestingly, Zeeman effect can produce FFLO superfluids condensed in higher bosonic Landau levels [21]. Typically, one Landau level is much more populated than the other ones, as shown in Fig.2 (a) and (c). Higher Landau levels can become favored at large h because they introduce additional vortex-antivortex pairs and therefore have larger regions of suppressed superfluidity where local spin polarization can be significant (see Fig.3). Structural transitions of the vortex lattice within the same Landau level are also found in the saddle point approximation, but quantum fluctuations of the order parameter generate familiar logarithmic “Coulomb” forces between vortices which generally act to stabilize the hexagonal Abrikosov lattice if it occurs in the lowest Landau level.

Finally, re-entrant superfluidity is reflected in the robust paired states which occur whenever the chemical potentials of the two atom species approach Landau levels. This is best demonstrated in Figure 4 which shows pairing free energy. As a result, a sequence of paired-state islands extends to arbitrary large h and survives at $h = 0$ regardless of how weak the attractive interactions between atoms are. The experimental consequences are discussed below.

The limitations of saddle point approximation (SPA) used to calculate the phase diagram in Fig.2 are discussed in [23]. An indicator for the quantitative applicability of SPA is $n\xi^2 \gg 1$, where n is atom density and $\xi \sim \sqrt{2\mu/m\Delta^2}$ the BCS coherence length (Δ is the spatially averaged BCS gap). This quantity grows rapidly with density or detuning as shown in Fig.5. The only qualitative shortcoming of SPA are the missed vortex liquid phases which are strongly correlated insulators of Cooper pairs with non-universal properties. These phases inevitably reside in the vicinity of all superfluid-insulator transitions shown in Fig.2 and grow at the expense of the vortex lattice regions, especially at the shown low densities when the number of atoms is not much larger than the number of vortices. The SPA still qualitatively captures the boundary between the paired correlated insulators and unpaired atomic quantum Hall states, as well as the bosonic Landau level occupation, and indirectly predicts the existence of exotic metallic and insulating spin-polarized vortex liquids. The first order transitions between vortex liquids at different densities and low temperatures are also expected to descend from the $T = 0$ “metal-insulator” transitions. We leave for future

research the more detailed studies of vortex liquids, which are potentially responsible for the unconventional properties of strongly correlated states in high-temperature superconductors [26].

The temperatures needed to explore the vortex-FFLO phase diagram in cold atom experiments should be below $T^* = k_B^{-1} \hbar \omega$, which is a few nK for typical $\omega \sim 2\pi \times 100$ Hz and within reach by evaporative cooling. The main difficulty would be working with a low enough density of trapped atoms, ideally not much larger than $n_0 = m\omega/(\pi\hbar) \sim 10^6 - 10^7$ cm⁻², the density for a single populated Landau level [27, 28]. However, the main features of the phase diagram in Fig.2 are expected to persist at much higher densities than shown, so that achieving the deep quantum limit may not be necessary. New experimental techniques may be able to overcome this difficulty in the future [29]. Fig.6 illustrates some possible experimental situations estimated from the grand-canonical results using local density approximation (LDA).

In this paper we analyzed a generic class of stable FFLO states in two-dimensions at zero temperature, whose influence is observable at finite temperatures. These states result from the combination of orbital and Zeeman effects which can be independently controlled in cold atom experiments. Further research, especially cold atom experiments near unitarity, could provide valuable answers to long standing questions about the stability of superconducting phases in large magnetic fields and the nature of strongly correlated normal states.

The mechanism for FFLO states described in this letter is perhaps the most generic and robust one. It requires a gap in the Bogoliubov quasiparticle spectrum smaller than the superfluid pairing scale. A vortex lattice is a natural agent to realize this situation due to the existence of sub-gap core states. One question is whether spontaneously developed spatial modulations in time-reversal symmetric conditions can produce such a quasiparticle band-structure that the band gap at the chemical potential is smaller than the superconducting gap. The answer to this question could be important for finding a stable traditional FFLO state.

I am very grateful to R.G.Hulet, Z.Hadzibabic and L.Baksmaty for useful comments and discussions. Numerical calculations were performed on Rice University computing clusters. This research was supported by W.M.Keck Program in Quantum Materials.

-
- [1] A.A.Abrikosov, *Zh.Eksp.Teor.Fiz.* **32**, 1442 (1957) [*Sov.Phys.JETP* **5**, 1174 (1957)].
 - [2] M.Rasolt, Z.Tešanović, *Rev.Mod.Phys.* **64**, 709 (1992).
 - [3] C.T.Rieck, K.Scharnberg, R.A.Klemm, *Physica C* **170**, 195 (1990).
 - [4] M.R.Norman, *Phys.Rev.B* **42**, 6762 (1990).
 - [5] P.Fulde, R.A.Ferrell, *Phys.Rev.* **135**, A550 (1964).
 - [6] A.I.Larkin, Yu.N.Ovchinnikov, *Zh.Eksp.Teor.Fiz.* **47**, 1136 (1964) [*Sov.Phys.JETP* **20**, 762 (1965)].
 - [7] U.Essmann, H.Träuble, *Phys.Lett.* **24A**, 526 (1967).
 - [8] I.Bloch, J.Dalibard, W.Zwenger, *Rev.Mod.Phys.* **80**, 885 (2008).
 - [9] M.W.Zwierlein, A.Schirotzek, C.H.Schunck, W.Ketterle, *Science* **311**, 492 (2006).
 - [10] Y.Shin, M.W.Zwierlein, C.H.Schunck, A.Schirotzek, W.Ketterle, *Phys.Rev.Lett.* **97**, 030401 (2006).
 - [11] G.B.Partridge, W.Li, R.I.Kamar, Y.Liao, R.G.Hulet, *Science* **311**, 503 (2006).
 - [12] G.B.Partridge *et al.*, *Phys.Rev.Lett.* **97**, 190407 (2006).
 - [13] R.Casalbuoni, G.Nardulli, *Rev.Mod.Phys.* **75**, 263 (2004).
 - [14] L.W.Gruenberg, L.Gunther, *Phys.Rev.Lett.* **16**, 996 (1966).
 - [15] A.I.Buzdin, J.P.Brison, *Phys.Lett.A* **218**, 359 (1996).
 - [16] K.Yang, A.H.MacDonald, *Phys.Rev.B* **70**, 094512 (2004).
 - [17] H.Shimahara, *J.Phys.Soc.Jpn.* **67**, 736 (1998).
 - [18] M.Houzet, A.Buzdin, L.Bulaevskii, M.Maley, *Phys.Rev.Lett.* **88**, 227001 (2002).
 - [19] H.Shimahara, *J.Phys.Soc.Jpn.* **67**, 1872 (1998).
 - [20] U.Klein, *Phys.Rev.B* **69**, 134518 (2004).
 - [21] Y.Matsuda, H.Shimahara, *J.Phys.Soc.Jpn.* **76**, 051005 (2007).
 - [22] N.R.Cooper, *Adv.Phys.* **57**, 539 (2008).
 - [23] P.Nikolić, *Phys.Rev.B* **79**, 144507 (2009).
 - [24] L.D.Landau, E.M.Lifshitz, *Course of Theoretical Physics, vol.5: Statistical Physics* (Butterworth-Heinemann, Oxford, 2000).
 - [25] S.Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 1999).
 - [26] L.Li, J.G.Checkelsky, S.Komiya, Y.Ando, N.P.Ong, *Nature Phys.* **3**, 311 (2007).

- [27] S.Stock, B.Battelier, V.Bretin, Z.Hadzibabic, J.Dalibard, *Laser Phys.Lett.* **2**, 275 (2005).
- [28] V.Schweikhard *et al.*, *Phys.Rev.Lett.* **92**, 040404 (2004).
- [29] K.J.Günter, M.Cheneau, T.Yefsah, S.P.Rath, J.Dalibard, *Phys.Rev.A* **79**, 011604(R) (2009).
- [30] G.B.Partridge, K.E.Strecker, R.I.Kamar, M.W.Jack, R.G.Hulet, *Phys.Rev.Lett.* **95**, 020404, (2005).
- [31] C.H.Schunck, Y.Shin, A.Schirotzek, M.W.Zwierlein, W.Ketterle, *Science* **316**, 867 (2007).




FIG. 1: Vortex core bands in relation to the superfluid gap near a vortex. The quasiparticle spectrum reduces to nearly-degenerate “Landau levels” at high energies, but resembles a band-structure shaped by the vortex lattice periodic potential at low energies, especially below the gap scale Δ_0 .




FIG. 2: Quantum phase diagrams ($T = 0$) of fast-rotating fermionic atoms near unitarity, showing spatially averaged density of Cooper pairs ρ_s and magnetization $m = n_\uparrow - n_\downarrow$: (a) ρ_s and (b) m at unitarity; (c) ρ_s and (d) m in the BCS limit at $\nu = 0.15$. The density plots of ρ_s are color-coded by the dominant Landau level (LL) n in the condensate, also indicated with numbers: red $n = 0$, green $n = 1$. The contour plots of m emphasize superfluid-insulator transitions with yellow lines (straight vertical segments are 2nd order, curved segments are 1st order), metal-insulator transitions within the superfluid phase with red lines (dotted in the ρ_s plots), and transitions between quantum Hall insulators (I) with thick-black lines. The superfluid phases (SF) host an Abrikosov vortex lattice, and may contain a coexisting metal (SF-M) or band-insulator (SF-I) of polarized unpaired fermions. The numbers in the phase labels indicate the highest populated quasiparticle band. Note that “spin-density” is saturated at an integer number of particles per flux quantum in all SF, SF-I and I phases, while it is not quantized in SF-M phases.




FIG. 3: Density plots of hexagonal vortex lattices for condensates in (a) $n = 0$, (b) $n = 1$, (c) $n = 2$, (d) $n = 3$, and (e) $n = 4$ Landau level. Brightness is normalized separately in each plot.




FIG. 4: Density plot of the pairing free energy density at (a) unitarity and (b) $\nu = 0.15$. Brightness indicates the amount of pairing.




FIG. 5: Parameter $n\xi^2$ at $h = 0$. Saddle point approximation is quantitatively justified if $n\xi^2 \gg 1$.




FIG. 6: Radial (r) profiles of Cooper pair density $\rho_s(r)$, total particle density $n(r)$ and polarization $p(r)$ as a function of total trap polarization $P = (N_\uparrow - N_\downarrow)/N$. The total particle number $N = N_\uparrow + N_\downarrow$ in the trap is $N = 3.5 \cdot 10^5$; dimensionless detuning is (a) $\nu = 0$, (b) $\nu = 0.15$. The total polarization P is indicated on the top. These plots illustrate sharp density features at very low temperatures. Starting from $P = 0$ the initial trend is that the superfluid core shrinks as the added “majority-spin” atoms are pushed to the trap boundary. However, the ring structure quickly acquires complexity from the Fig.2: moving from $r = 0$ toward the trap edge follows a trajectory of reducing μ while keeping h constant in Fig.2. Characteristic superfluid rings separated by normal regions are a signature of re-entrant superfluidity, and polarization can be finite in a superfluid ring (FFLO states, or SF-I and SF-M in Fig.2). The distinction between SF-I and SF-M phases can be made by whether $p(r)$ is a quantized constant or smoothly varying respectively. The SF regions replaced by vortex liquids are expected to have strong pairing fluctuations [30, 31]. A currently experimentally unreachable trap harmonic frequency $\omega_\perp = (1 + 5 \cdot 10^{-5})\omega$ was chosen to justify local density approximation, and the shown spatial region is $r < 500\sqrt{\hbar/(2m\omega)}$. All other parameters are comparable with already accomplished experimental setups [27, 28].